

AD-A193 117


HELICAL LATTICE VIBRATIONAL MODES IN DNA(U) PURDUE UNIV 1/1
LAFAYETTE IND DEPT OF PHYSICS V V PRADHU ET AL.
10 MAR 89 N00014-86-K-0252

UNCLASSIFIED

F/G 6/2

NL





MICROCOPY RESOLUTION TEST CHART
 NATIONAL BUREAU OF STANDARDS-1963-A

mm

FILE COPY

REPORT DOCUMENTATION PAGE

2

AD-A193 117

1. AUTHORING ORGANIZATION NAME(S)		2. DISTRIBUTION STATEMENT OF REPORT	
Purdue Research Foundation		Distribution Unlimited	
3. MONITORING ORGANIZATION REPORT NUMBER(S)		4. NAME OF MONITORING ORGANIZATION	
NA		Technology Applications Programs	
5a. NAME OF FUNDING SPONSORING ORGANIZATION	5b. OFFICE SYMBOL (If applicable)	7a. NAME OF MONITORING ORGANIZATION	
Purdue Research Foundation	NA	Technology Applications Programs	
6a. ADDRESS (City, State and ZIP Code)		7b. ADDRESS (City, State and ZIP Code)	
W. Lafayette, IN 47907		OSD/SD10/TA Washington, D.C. 20301-7100	
8a. NAME OF FUNDING SPONSORING ORGANIZATION	8b. OFFICE SYMBOL (If applicable)	9. ACQUISITION INSTRUMENT IDENTIFICATION NUMBER	
Strategic Defense Initiative Organization		N00014-86-K-0252	
5c. ADDRESS (City, State and ZIP Code)		10. SOURCE OF FUNDING NUMBERS	
Washington, D.C. 20301-7100		PROGRAM ELEMENT NO 63222C	
		PROJECT NO	TASK NO
			WORK UNIT ACCESSION NO
11. TITLE (Include Security Classification)			
Helical Lattice Vibrational Modes in DNA			
12. PERSONAL AUTHOR(S)			
V.V. Prabhu, W.K. Schroll, L.L. Van Zandt, E.W. Prohofsky			
13a. TYPE OF REPORT	13b. TIME COVERED	14. DATE OF REPORT (Year, Month, Day)	15. PAGE COUNT
Technical	FROM 4/81 TO 3/88	1988 - 3-10	
16. SUPPLEMENTARY NOTES			
17. COSAT CODES		18. SUBJECT TERMS (Continue on reverse if necessary and identify by block number)	
FIELD	GROUP	SUB GROUP	
19. ABSTRACT (Continue on reverse if necessary and identify by block number)			
<p>We show that the acoustic branches calculated by our helical lattice dynamics calculations can match the observed inelastic neutron scattering results. The fit in the neutron sub Terahertz region is consistent with our fit to Brillouin results in the Ggahertz region when the frequency dependence of the dielectric constant of water is taken into account.</p>			
<div style="text-align: right;"> <p>DTIC ELECTE</p> <p>S APR 11 1988 D</p> <p>H</p> </div>			
20. DISTRIBUTION AVAILABILITY OF ABSTRACT		21. ABSTRACT SECURITY CLASSIFICATION	
<input type="checkbox"/> UNCLASSIFIED/UNLIMITED <input type="checkbox"/> SAME AS REF <input type="checkbox"/> DTIC USERS		U	
22a. NAME OF RESPONSIBLE INDIVIDUAL		22b. TELEPHONE (Include Area Code)	22c. OFFICE SYMBOL
Charles L. Houston III		(202)693-1556	OSD/SD10/TA

DISTRIBUTION STATEMENT A

Approved for public release:
Distribution Unlimited

Helical Lattice Vibrational Modes in DNA

A recent letter by Grimm et al.¹ reported the observation of umklapp-phonon scattering at sub teraHertz frequencies in DNA by inelastic neutron scattering. They found, that the nearly linear dispersion of their results did not extrapolate to zero frequency at zero wavevector as a simple acoustic mode should and that the data could be best fit by a dispersion relation $\omega^2 = \omega_0^2 + v^2(\Delta Q)^2$ where $\omega_0/2\pi$ was 0.4 THz. This is in disagreement with two earlier observations^{2,3} carried out by Brillouin scattering, between 5-20 GHz, where the acoustic mode is seen to behave normally and extrapolate to zero frequency at zone center. We point out, that a proper helical lattice dynamics analysis of vibrational modes of the DNA helix resolves this apparent inconsistency, and that the inelastic neutron data as well as the Brillouin data are in excellent quantitative agreement with such calculations.

The lowest optical mode at zone center in Fig. 1 has been observed experimentally⁴, as well as other optical modes calculated by the lattice theory.⁵ As seen in Fig. 1 the lowest longitudinal acoustic branch does extrapolate linearly to zero frequency at zone center. The Brillouin observations in the GHz region fall on this branch. The heavy lines indicate the continuation of the compressional character of the phonons on the successive branches of the dispersion plot of Fig. 1. The experimental resolution is such that the splitting of the branches near the crossover cannot be seen. We note that a feature exists in the experimental data near the second from bottom optic-acoustic crossover that is consistent with the predicted region of crossover.

The value of the acoustic velocity from this calculation is 4.5 km/s. To compare with the Brillouin work in the GHz region, we changed the long range water dielectric to 45 (from 9) and calculated an acoustic velocity of

3.1 km/s. This is about what would be expected for a helix unburdened by water of hydration, based on measurements by Lee et al.⁶

This work is supported by NIH Grant GM 24443 and ONR grants N00014-86-K-0252 and N00014-87-K-0162.

V. V. Prabhu
W. K. Schroll
L. L. Van Zandt
E. W. Prohofsky

Department of Physics
Purdue University
West Lafayette, IN 47907

PACS numbers: 87.15.-v, 63.20.Dj


Accession For	
NTIS GRA&I	<input checked="checked" type="checkbox"/>
DTIC TAB	<input type="checkbox"/>
Unannounced	<input type="checkbox"/>
Justification	
By	
Distribution/	
Availability Codes	
Dist	Avail and/or Special
A-1	For INSPE

References

1. H. Grimm, H. Stiller, C. F. Majkrzak, A. Rupprecht, U. Dahlborg, Phys. Rev. Lett. 59, 1780 (1987).
2. G. Maret, R. Oldenbourg, G. Winterling, K. Dransfeld and A. Rupprecht, Colloid Polym. Sci. 257, 1017-1020, (1979)
3. S. M. Lindsay and J. Powell, Structure and Dynamics - Nucleic Acids and Proteins. E. Clementi and R. H. Sarma, Eds., Adenine Press, New York, pp. 241-260.
4. S. M. Lindsay, J. Powell, E. W. Prohofsky, and K. V. Devi-Prasad. Structure and Motion: Membranes, Nucleic Acids and Proteins, E. Clementi, G. Corongiu, M. H. Sarma, and R. H. Sarma, Eds. Adenine Press, New York (1985) pp. 531-551.
5. K. V. Devi-Prasad and E. W. Prohofsky, Biopolymers 23, 1795-1798 (1984).
6. S. A. Lee. S. M. Lindsay, J. W. Powell, T. Weidlich, N. J. Tao, G. D. Lewen and A. Rupprecht, Biopolymers 26, 1637-1665 (1987).

Figure Captions

Fig. 1 The continuous lines are our helical lattice dispersion calculations of the phonon spectrum of DNA. The θ axis is displayed in degrees of phase shift from one unit cell of two DNA base pairs to the next. The central row of stars show the reported frequencies from the neutron data and the outer rows of dashes show the reported FWHM line width. The heavy lines indicate the continuation of the compressional character of the phonons on the successive branches of the dispersion plot.



END

DATE

FILMED

DTIC

JULY 88